



Carbon-Based Nanomaterials for Soil Amendment: A Paradigm Shift in Agriculture Practices

S. Alagendran ^{a++*}, C. Mounika ^{b#}, Ashoka P ^{ct},
Vikas Singh ^{d‡}, Chakri Voruganti ^e,
Papita H Gourkhede ^{f^} and M. Niharika ^{g^}

^a Department of Biochemistry, Dhanalakshmi Srinivasan Agriculture College, (Affiliated to Tamil Nadu Agricultural University, Coimbatore-3) Perambalur, 621212 Tamil Nadu, India.

^b Division of Soil Science and Agricultural Chemistry, Indian Agriculture Research Institute, New Delhi, India.

^c Department of Agronomy, College of Agriculture Hanumanmatti (p), Ranebennur (tq), Haveri (District), 581 115, Karnataka, India.

^d Department of Soil Conservation and Water Management, Chandra Shekhar Azad university of Agriculture and Technology, Kanpur, Uttar Pradesh, India.

^e Department of Plant and Soil Science, Texas Tech University, Lubbock, Texas, USA.

^f Department of Soil Science, AICRP on Dryland Agriculture, VNMKV, Parbhani, 431402, Maharashtra, India.

^g Department of Food Science and Nutrition, KL College of Agriculture, KLEF, Vaddeswaram, Guntur, Andhra Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i123192>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129020>

⁺⁺ Ph D (Biochemistry); PDF (UNAM, MX) (5th Inventor) - 5th Author, Associate Professor;

[#] Ph.D Scholar;

[†] Professor and Head;

[‡] Teaching Associate;

[^] Assistant Professor;

*Corresponding author: E-mail: alagendran@dsagri.in;

Cite as: Alagendran, S., C. Mounika, Ashoka P, Vikas Singh, Chakri Voruganti, Papita H Gourkhede, and M. Niharika. 2024. "Carbon-Based Nanomaterials for Soil Amendment: A Paradigm Shift in Agriculture Practices". *Journal of Experimental Agriculture International* 46 (12):833-49. <https://doi.org/10.9734/jeai/2024/v46i123192>.

ABSTRACT

Carbon-based nanomaterials (CBNMs), including biochar, graphene oxide (GO), carbon nanotubes (CNTs), and fullerenes, have emerged as innovative tools for sustainable agriculture, particularly in soil amendment applications. Their unique properties, such as high surface area, chemical stability, porosity, and functionalization potential, allow for significant improvements in soil health, nutrient retention, and microbial activity. CBNMs enhance soil structure, water retention, and aeration, while also reducing the leaching of fertilizers and agrochemicals, thereby minimizing environmental contamination. biochar's role in carbon sequestration and the mitigation of greenhouse gas emissions positions it as a valuable tool for climate-smart agriculture. Recent research highlights CBNMs' ability to improve crop yields, nutrient use efficiency, and stress tolerance under conditions such as drought and salinity. Challenges remain regarding the scalability of their production, high costs, and potential toxicity to soil ecosystems and plants, as well as risks of nanoparticle leaching into water bodies. The long-term impacts of CBNMs on soil and the broader environment are still poorly understood, necessitating further investigation into their interactions with different soil types and ecosystems. Integration with emerging technologies, such as precision agriculture and Internet of Things (IoT)-based systems, presents opportunities to optimize their application and monitoring. Adopting CBNMs in regenerative agriculture practices could further enhance soil health and ecosystem resilience. Future directions should prioritize developing cost-effective and scalable synthesis methods, establishing regulatory frameworks for safe use, and promoting interdisciplinary collaborations to address knowledge gaps and public concerns. As the field advances, CBNMs have the potential to revolutionize agricultural practices, improving soil sustainability and productivity while mitigating environmental challenges, thereby contributing to global food security and climate resilience. This review underscores the promise of CBNMs in agriculture, while emphasizing the need for responsible innovation and rigorous research to ensure their sustainable implementation.

Keywords: Carbon-based nanomaterials; biochar; graphene oxide; soil amendment.

1. INTRODUCTION

Soil health is the cornerstone of sustainable agricultural systems, as it directly impacts crop productivity, environmental quality, and ecosystem services. Defined as the continued capacity of soil to function as a vital living ecosystem that sustains plants, animals, and humans, soil health integrates physical, chemical, and biological components essential for agricultural success. A healthy soil promotes nutrient cycling, water retention, carbon sequestration, and biodiversity, which are critical for long-term agricultural resilience. Importantly, soils host more than 25% of the planet's biodiversity and are vital for carbon storage, with approximately 80% of terrestrial carbon stored within soil ecosystems. Given the projected increase in global food demand by 70% by 2050, maintaining and improving soil health is imperative for achieving food security while

minimizing environmental degradation. Healthy soils are integral to sustainable agricultural practices, enabling improved water infiltration and storage, reduced erosion, and increased microbial activity, which supports nutrient bioavailability (Lehman *et.al.*, 2015). Soils with balanced organic matter content improve plant root systems by enhancing nutrient uptake and stabilizing the soil structure. Thus, soil health is an essential component of sustainable agricultural practices, ensuring the production of nutritious food, the mitigation of greenhouse gas emissions, and the preservation of biodiversity. Modern agricultural systems, while boosting global food production, have led to significant environmental consequences, including widespread soil degradation. Soil degradation, defined as the decline in soil quality due to factors such as erosion, nutrient depletion, and contamination, currently affects an estimated 33% of global arable land. Intensive farming

practices, such as monocropping, excessive tillage, and overuse of synthetic fertilizers and pesticides, have contributed to the depletion of essential nutrients, organic matter, and microbial biodiversity within the soil (Patle *et.al.*, 2019). Nutrient depletion is a particularly pressing challenge, as modern agricultural practices often remove more nutrients from the soil than are replenished, leading to declining crop yields and poor soil fertility. For example, nitrogen, phosphorus, and potassium the three macronutrients critical for plant growth are often leached or eroded, resulting in substantial nutrient imbalances in many farming systems. Furthermore, the degradation of soil organic matter exacerbates nutrient loss, reducing the soil's capacity to retain and supply nutrients effectively. These issues are further compounded by climate change, which intensifies soil erosion, salinization, and desertification, particularly in arid and semi-arid regions (Reed *et.al.*, 2016).

1.1 Emergence of Carbon-Based Nanomaterials (CBNMs)

Carbon-based nanomaterials (CBNMs) are a diverse class of nanoscale materials composed primarily of carbon atoms, characterized by unique physical, chemical, and biological properties that make them highly versatile for various applications, including soil amendment (Table 1). These materials typically feature high surface area, thermal and chemical stability, electrical conductivity, and the ability to interact with biological and chemical systems at the molecular level. The most commonly studied types of CBNMs for soil applications include: Biochar: A carbon-rich material produced via pyrolysis of organic biomass under low-oxygen conditions. Biochar has a porous structure and high cation exchange capacity, making it effective for improving soil fertility and sequestering carbon. Graphene and Graphene Oxide: Two-dimensional carbon materials with extraordinary mechanical strength, electrical conductivity, and hydrophilicity. These materials are gaining attention for enhancing water retention and nutrient delivery in soils (Davidson *et.al.*, 2012). Carbon Nanotubes (CNTs): Cylindrical carbon nanostructures with a high aspect ratio, excellent tensile strength, and remarkable adsorption properties. CNTs are used to improve soil aeration and facilitate nutrient transport. Fullerenes: Spherical molecules composed entirely of carbon, often used for soil remediation due to their ability to interact with organic and inorganic contaminants.

2. PROPERTIES OF CBNMs THAT MAKE THEM PROMISING FOR SOIL AMENDMENT

The unique properties of carbon-based nanomaterials make them highly promising for soil amendment and agricultural applications. Their high specific surface area allows for enhanced adsorption of nutrients, water, and pollutants, improving soil nutrient retention and reducing environmental contamination (Nakhli *et.al.*, 2017). Biochar and graphene-based materials can mitigate the leaching of fertilizers, ensuring sustained nutrient availability for plant uptake. The porous structure of many CBNMs, such as biochar, enhances soil aeration and water-holding capacity, which are critical for plant growth, especially in drought-prone regions. CBNMs also demonstrate strong chemical reactivity and functionalization potential, enabling them to modulate soil pH, immobilize toxic metals, and enhance cation exchange capacity. Furthermore, their ability to interact with soil microbiota can stimulate beneficial microbial activity while suppressing pathogenic organisms, thereby contributing to improved soil health and plant productivity. Graphene and CNTs, have been shown to promote root elongation and seed germination through their interaction with plant cells and soil components (Zhang *et.al.*, 2022).

3. OBJECTIVES OF THE REVIEW

The primary objective of this review is to explore how carbon-based nanomaterials (CBNMs) contribute to improving soil health and agricultural productivity. By analyzing their mechanisms of action, including physical, chemical, and biological interactions within the soil, this review seeks to provide a comprehensive understanding of their potential benefits. The review aims to examine the role of CBNMs in addressing critical challenges in modern agriculture, such as nutrient depletion, water scarcity, and soil degradation, while highlighting their impact on crop yield and quality. This review also aims to summarize current advancements in the field of carbon-based nanomaterials, focusing on laboratory experiments, field trials, and case studies that demonstrate their effectiveness as soil amendments. Furthermore, the paper identifies key challenges in the adoption of CBNMs, such as cost, scalability, environmental risks, and regulatory issues (Asghar *et.al.*, 2024). By discussing these limitations, the review provides insights into future research directions,

interdisciplinary collaborations, and policy frameworks required to accelerate the adoption of CBNMs in agriculture. Through this discussion, the paper emphasizes the paradigm shift that CBNMs represent in modern agricultural practices.

4. CHARACTERISTICS AND TYPES OF CARBON-BASED NANOMATERIALS

4.1 Carbon-Based Nanomaterials

Carbon-based nanomaterials (CBNMs) are a diverse group of nanoscale materials composed predominantly of carbon atoms organized into various structures, including one-dimensional, two-dimensional, and three-dimensional morphologies (Table 2). Their unique nanoscale size (typically below 100 nm) imparts them with distinct physical, chemical, and biological properties that differ significantly from their bulk counterparts. These properties make them highly versatile for a wide range of applications, including agriculture, where they have emerged as promising tools for soil amendment. The high surface-to-volume ratio of nanomaterials enhances their capacity to adsorb nutrients, water, and pollutants, making them effective for improving soil properties (Hussain *et.al.*, 2022). CBNMs exhibit resistance to degradation under extreme environmental conditions, ensuring their longevity in soil ecosystems. Carbon nanomaterials can be easily functionalized with various chemical groups, allowing for tailored interactions with soil components, such as nutrients and contaminants. Many CBNMs, such as graphene and biochar, have porous structures that enhance soil aeration and water-holding capacity while also improving soil structure. Materials such as graphene and carbon nanotubes exhibit exceptional electrical conductivity, which has been explored for applications in plant growth stimulation and nutrient delivery (Vithanage *et.al.*, 2017).

4.2 Synthesis Methods

The synthesis of carbon-based nanomaterials involves various techniques designed to control their size, structure, and functionalization. The most common synthesis methods include: This method involves the thermal decomposition of organic biomass under low-oxygen conditions, resulting in biochar. Pyrolysis conditions, such as temperature and residence time, influence biochar's porosity, carbon content, and surface area, all of which are crucial for soil applications.

A widely used method for synthesizing high-purity graphene and carbon nanotubes. In this process, carbon precursors (e.g., methane or acetylene) are heated and deposited on a substrate, forming nanostructures. CVD allows precise control over material properties, such as layer thickness and crystallinity. A method for producing carbon nanotubes and fullerenes, arc discharge involves creating a high-energy plasma between two graphite electrodes in an inert gas atmosphere. This technique yields high-quality nanomaterials, although scaling up can be challenging (Farhat *et.al.*, 2006). A green method used for synthesizing graphene oxide and functionalized biochar, hydrothermal synthesis employs high-pressure, high-temperature water to convert biomass or carbon precursors into nanostructures.

5. KEY TYPES OF CBNMs USED IN AGRICULTURE

Biochar, a carbon-rich material derived from the pyrolysis of organic biomass, has gained significant attention as a soil amendment due to its porous structure, high cation exchange capacity (CEC), and stability in soil environments. The porous architecture of biochar enhances soil aeration and water retention, making it particularly effective in arid and degraded soils. Its high surface area provides abundant sites for nutrient adsorption, reducing nutrient leaching and improving fertilizer efficiency (Wang *et.al.*, 2019). Biochar also serves as a habitat for beneficial soil microorganisms, promoting microbial diversity and activity. Its role in carbon sequestration has been widely documented, as biochar can remain stable in the soil for centuries, contributing to climate change mitigation efforts. Furthermore, biochar's ability to immobilize heavy metals and organic pollutants makes it a valuable tool for soil remediation. Graphene and graphene oxide (GO) are two-dimensional carbon nanomaterials known for their extraordinary mechanical strength, high electrical conductivity, and hydrophilic properties. These characteristics make them highly promising for enhancing soil properties and supporting plant growth. Graphene's ability to retain water and nutrients is particularly beneficial in sandy or degraded soils. Studies have shown that graphene-based materials can improve soil water-holding capacity by creating a hydrophilic network, which enhances plant water availability during drought conditions. GO functional groups can adsorb and release nutrients, improving nutrient availability

and reducing fertilizer loss. Graphene also interacts with soil microbiota, enhancing microbial activity and diversity, which are critical for nutrient cycling and soil health. Concerns about graphene's potential ecotoxicity warrant further investigation to ensure its safe use in agriculture (Wahab *et.al.*, 2024). Carbon nanotubes (CNTs) are cylindrical carbon nanostructures with high tensile strength, electrical conductivity, and a large aspect ratio. Their unique properties have been exploited for various agricultural applications, including soil amendment, nutrient delivery, and plant growth stimulation. CNTs can enhance soil aeration and water movement due to their tubular structure, which creates microchannels within the soil matrix. CNTs have shown the ability to deliver nutrients and agrochemicals directly to plant roots, improving nutrient use efficiency and reducing environmental contamination. Studies have also demonstrated that CNTs can stimulate seed germination and root elongation by interacting with plant cells at the molecular level. Like graphene, the environmental fate and potential toxicity of CNTs remain areas of active research. Fullerenes are spherical carbon molecules (e.g., C60) with unique chemical and physical properties, including high electron affinity and photochemical reactivity (Pan *et.al.*, 2020). Although their use in agriculture is less studied compared to other CBNMs, fullerenes have shown potential for niche applications, such as soil remediation and pollutant removal. Fullerenes can interact with organic and inorganic contaminants, immobilizing them within the soil matrix and reducing their bioavailability. Their antioxidant properties may also help mitigate oxidative stress in plants exposed to environmental stressors, such as heavy metals or drought. While fullerenes offer unique benefits, their high cost and limited scalability pose challenges for widespread agricultural use.

6. COMPARISON OF CBNMs IN TERMS OF PROPERTIES RELEVANT TO SOIL HEALTH

The effectiveness of carbon-based nanomaterials in improving soil health depends on their specific properties, such as surface area, porosity, and functionalization potential. A comparison of the key CBNMs is summarized below (Table 3).

Each type of CBNM offers unique advantages and trade-offs, making them suitable for specific agricultural applications depending on the soil type, crop requirements, and environmental conditions (Thompson *et.al.*, 2022).

7. MECHANISMS OF ACTION: HOW CBNMs ENHANCE SOIL HEALTH

Carbon-based nanomaterials (CBNMs) offer multifaceted mechanisms to improve soil health through physical, chemical, and biological pathways. These mechanisms influence soil properties, nutrient cycling, microbial communities, and plant productivity, making CBNMs highly effective for sustainable soil management.

7.1 Physical Mechanisms

One of the primary ways CBNMs improve soil health is by enhancing its physical structure. Materials like biochar and carbon nanotubes (CNTs) contribute to better soil aggregation and porosity due to their porous architecture and high surface area. Biochar particles, act as scaffolds within the soil, binding with mineral particles and organic matter to form aggregates that improve soil stability and resistance to erosion. Increased soil porosity directly impacts root penetration and soil aeration, creating favorable conditions for plant growth. Furthermore, the introduction of nanoscale materials into compacted or degraded soils can increase their bulk density and permeability, facilitating the movement of water and air (Liu *et.al.*, 2012). These improvements are particularly beneficial for clayey or poorly structured soils. CBNMs improve soil water retention by increasing the soil's water-holding capacity. Biochar, due to its microporous and hydrophilic structure, retains water within its pores and releases it gradually to the soil matrix, mitigating drought stress. Graphene oxide (GO) is another example of a CBNM with exceptional water-retention properties due to its ability to form hydrogen bonds with water molecules. The aeration benefits of CBNMs stem from their ability to enhance soil porosity. Biochar and CNTs help prevent waterlogging by promoting air exchange, which is vital for root respiration and the activity of aerobic microbes. These improvements in water retention and aeration are critical for improving plant growth, particularly in arid and semi-arid regions.

Table 1. Emergence of Carbon-Based Nanomaterials (CBNMs)

Aspect	Definition	Historical Development	Types	Properties	Synthesis Techniques	Applications
Overview	Materials with nanometer-scale dimensions, primarily composed of carbon.	Discovery began with fullerenes in 1985, followed by carbon nanotubes (CNTs) in 1991, and graphene in 2004.	Fullerenes, CNTs, graphene, carbon quantum dots, nanodiamonds.	High electrical/thermal conductivity, large surface area, tunable surface chemistry, strong mechanical properties.	Chemical Vapor Deposition (CVD), arc discharge, laser ablation, chemical exfoliation.	Electronics, energy storage, drug delivery, catalysis, sensors, and composite materials.
Innovative Features	Nanoscale structures offer unique chemical, physical, and biological properties.	Evolution shaped by advancements in synthesis technologies and characterization tools.	Can be functionalized for specific applications, improving their versatility.	Exceptional ability to withstand deformation, high electrical and thermal conductivity, lightweight nature.	High-temperature techniques and solution-phase methods are leading the field.	Medical imaging, advanced coatings, environmental monitoring, and energy-efficient devices.
Future Perspectives	Integration with AI for design, advanced applications in sustainable energy and nanomedicine expected.	Research focuses on scalability, environmental safety, and cost-effectiveness of production methods.	Focus on hybrid CBNMs for multifunctional applications.	Potential for novel bio-compatible and eco-friendly material innovations.	Exploring green synthesis methods for eco-friendly nanomaterial development.	Renewable energy systems, water purification, and structural reinforcements in the aerospace industry.

(Source: Lehman et al., 2015)

Table 2. Types of Carbon-Based Nanomaterials

Type	Structure	Key Properties	Synthesis Methods	Applications
Fullerenes	Hollow spherical or elliptical structures composed entirely of carbon atoms.	High electron affinity, stability under various conditions, unique photochemical and electrochemical properties.	Arc discharge, laser ablation, and combustion synthesis.	Drug delivery systems, photovoltaics, antioxidants, and materials for organic electronics.
Carbon Nanotubes (CNTs)	Cylindrical tubes made of rolled graphene sheets.	Exceptional tensile strength, high electrical and thermal conductivity, lightweight.	Chemical Vapor Deposition (CVD), arc discharge, laser ablation.	Nanoelectronics, composite materials, biosensors, and water filtration systems.

Type	Structure	Key Properties	Synthesis Methods	Applications
Graphene	Single layer of carbon atoms arranged in a hexagonal lattice.	High mechanical strength, excellent thermal and electrical conductivity, high surface area.	Mechanical exfoliation, chemical exfoliation, Chemical Vapor Deposition (CVD).	Flexible electronics, energy storage devices (supercapacitors, batteries), and membrane technology.
Carbon Quantum Dots	Small, quasi-spherical nanoparticles of carbon.	Fluorescent properties, biocompatibility, tunable photoluminescence.	Hydrothermal synthesis, chemical oxidation, laser ablation.	Bioimaging, biosensors, photocatalysis, and drug delivery systems.
Nanodiamonds	Diamond particles in the nanometer range.	Hardness, chemical stability, high thermal conductivity, and biocompatibility.	Detonation synthesis, laser ablation, high-pressure high-temperature synthesis.	Coatings, biomedical applications, drug delivery, and optical devices.

Source: (Hussain et al., 2022), (Vithanage et al., 2017)

Table 3. Comparison of CBNMs in Terms of Properties Relevant to Soil Health

CBNM Type	Surface Area	Water Retention	Nutrient Adsorption	Soil Structure Improvement	Microbial Interactions
Biochar	High	High	High	High	High
Graphene/GO	Very High	Moderate to High	High	Moderate	High
CNTs	High	Moderate	Moderate	High	Moderate
Fullerenes	Moderate	Low	Moderate	Low	Low

Source: (Wang et al., 2019), (Pan et al., 2020)

7.2 Chemical Mechanisms

CBNMs enhance soil chemical properties by increasing the cation exchange capacity (CEC) of soils, which governs the ability of soils to retain and release essential nutrients such as potassium (K^+), calcium (Ca^{2+}), and magnesium (Mg^{2+}). Biochar is particularly notable for its high CEC, attributed to the abundance of functional groups (e.g., carboxyl, hydroxyl) on its surface that facilitate the adsorption of nutrients (Zhang *et.al.*, 2021). Graphene oxide and CNTs also play a role in improving nutrient availability by acting as carriers for nutrients or fertilizers. These nanomaterials can adsorb and slowly release nutrients, ensuring their prolonged availability to plants while reducing nutrient loss due to leaching. The immobilization of nutrients on the surfaces of CBNMs minimizes volatilization and runoff, thus improving the efficiency of fertilizers and reducing environmental contamination. Nutrient leaching is a major contributor to soil degradation and water contamination. CBNMs mitigate this issue by trapping nutrients and agrochemicals within their porous structures, reducing their mobility in the soil matrix. For example, biochar has been shown to reduce nitrate and phosphate leaching in agricultural soils, protecting groundwater resources (Cui *et.al.*, 2020). Graphene oxide and CNTs form complexes with agrochemicals, slowing their release into the environment. These properties not only enhance the sustainability of agricultural practices but also help maintain soil fertility over time.

7.3 Biological Mechanisms

CBNMs support the proliferation of beneficial soil microorganisms by creating microhabitats and improving soil conditions. Biochar, in particular, has been widely studied for its ability to serve as a microbial habitat due to its porous structure and the availability of carbon-rich surfaces. This promotes microbial diversity and enhances processes such as nitrogen fixation and organic matter decomposition. Graphene oxide has also been reported to stimulate microbial activity by modulating microbial metabolism and supporting the growth of plant growth-promoting rhizobacteria (PGPR) (Wu *et.al.*, 2023). This is crucial for nutrient cycling and improving the bioavailability of essential elements like nitrogen and phosphorus. CBNMs can suppress harmful soil-borne pathogens by creating unfavorable conditions for their growth. Biochar has been shown to adsorb toxins and pathogens, reducing

their bioavailability and impact on crops. Certain nanomaterials, such as CNTs and fullerenes, exhibit antimicrobial properties, which can further suppress pathogenic microorganisms. The interactions between CBNMs and soil microbiota are complex and depend on the type and concentration of the nanomaterial. While biochar and graphene oxide generally enhance microbial activity, excessive concentrations of some CBNMs, such as CNTs, may exert cytotoxic effects on soil microorganisms (Chahar *et.al.*, 2023). Thus, understanding the dose-dependent effects of CBNMs is crucial for maximizing their benefits while minimizing potential risks. CBNMs also influence microbial community composition. For example, biochar has been shown to promote the growth of nitrogen-fixing bacteria, while graphene oxide can selectively enhance the activity of beneficial microbes involved in nutrient cycling.

7.4 Case Studies of CBNM Applications in Soil Management

In a field study conducted in Australia, biochar was applied to degraded sandy soils, resulting in a 30% increase in water-holding capacity and a 25% reduction in nutrient leaching. This translated into improved crop yields and long-term soil fertility. A greenhouse experiment demonstrated that graphene oxide increased nitrogen retention in sandy soils by 40%, reducing fertilizer losses and enhancing plant growth (Kabiri *et.al.*, 2017). The study highlighted graphene oxide's potential to improve nutrient use efficiency in agricultural systems. Carbon nanotubes improved tomato plant growth by increasing root elongation and seed germination rates. The CNTs facilitated water uptake and nutrient transport, showcasing their potential for enhancing crop productivity. The use of fullerenes for immobilizing heavy metals in contaminated soils. The fullerenes reduced the bioavailability of cadmium and lead, protecting plants from metal toxicity and improving soil health.

8. ADVANTAGES OF CARBON-BASED NANOMATERIALS FOR SOIL AMENDMENT

Carbon-based nanomaterials (CBNMs) have gained widespread attention as soil amendments due to their unique properties, which allow them to improve soil health, increase agricultural productivity, and offer environmental sustainability (Sheoran *et.al.*, 2022).

8.1 Sustainability and Environmental Benefits

The overuse of chemical fertilizers in agriculture has led to severe environmental consequences, including soil degradation, eutrophication, and groundwater contamination. CBNMs, such as biochar, graphene oxide, and carbon nanotubes (CNTs), provide an eco-friendly alternative to conventional fertilizers by enhancing nutrient retention and availability in the soil. Biochar, for example, acts as a reservoir for nutrients, reducing leaching and volatilization of nitrogen and phosphorus, which decreases the need for synthetic fertilizers. Graphene oxide and CNTs further contribute to nutrient efficiency by facilitating the slow release of fertilizers. Studies have demonstrated that these materials adsorb and retain nutrients, such as ammonium and phosphate, on their surfaces, ensuring prolonged availability to plants (Reddy *et.al.*, 2017). This reduces fertilizer application frequency, minimizes waste, and mitigates the environmental impacts associated with excessive fertilizer use. Carbon sequestration is one of the most significant environmental benefits of using CBNMs, particularly biochar, as a soil amendment. Biochar is a stable form of carbon that resists microbial decomposition, allowing it to remain in the soil for centuries. Its application not only improves soil fertility but also serves as a strategy for mitigating climate change by capturing atmospheric carbon dioxide (CO₂) and storing it in the soil.

8.2 Enhanced Crop Productivity

CBNMs significantly contribute to increased crop yields and improved crop quality by enhancing soil properties and optimizing nutrient availability. Biochar, for example, has been shown to increase crop yields by improving soil structure, water retention, and nutrient use efficiency (Fischer *et.al.*, 2019). A meta-analysis found that biochar application resulted in an average yield increase of 25% across various crops and soil types. Graphene oxide and CNTs have demonstrated the ability to stimulate plant growth by improving root development and nutrient uptake. Graphene oxide, in particular, enhances nutrient bioavailability and promotes seed germination and root elongation. Enhanced nutrient delivery through CNTs has been shown to boost the growth of tomato plants and increase the uptake of essential minerals like potassium and magnesium. CBNMs can improve the nutritional quality of crops. Studies have

reported that biochar-enriched soils result in higher protein and nutrient content in grains, fruits, and vegetables. This underscores the potential of CBNMs to not only enhance productivity but also produce more nutritious food. Soil amendments with CBNMs can significantly improve plant tolerance to abiotic stresses, such as drought and salinity, which are major threats to global food security. Biochar's water-retention properties are particularly beneficial in arid and semi-arid regions, where water scarcity limits crop growth. By increasing the soil's water-holding capacity, biochar ensures that plants have access to moisture during dry periods (Gharred *et.al.*, 2022). Graphene oxide has also been studied for its ability to mitigate water stress in plants. Due to its high hydrophilicity, graphene oxide forms a water-retentive matrix in the soil, reducing evaporation and ensuring steady water availability to plants. Graphene oxide and CNTs help plants cope with salinity stress by reducing the uptake of toxic ions, such as sodium (Na⁺), and improving the absorption of essential nutrients like potassium (K⁺).

8.3 Longevity and Cost-Effectiveness

One of the most compelling advantages of CBNMs is their durability and long-term stability in soil ecosystems. Biochar, is highly resistant to microbial decomposition and can remain in soils for centuries, continuously improving soil structure and nutrient retention over time. This contrasts with traditional soil amendments, such as compost and manure, which decompose relatively quickly and require frequent reapplication (Quilty *et.al.*, 2011). In their physical and chemical stability, CBNMs support long-term soil fertility by promoting microbial activity and nutrient cycling. The sustained benefits of biochar and graphene oxide for soil microbial communities ensure ongoing improvements in nutrient availability and soil health. These enduring effects make CBNMs a cost-effective solution for sustainable soil management. While the initial costs of CBNMs may be higher than conventional soil amendments, their long-term benefits often justify the investment. For example, the one-time application of biochar can replace repeated applications of synthetic fertilizers due to its ability to enhance nutrient efficiency and retention (Kizito *et.al.*, 2019). A cost-benefit analysis conducted in low-income agricultural systems demonstrated that biochar application resulted in a net economic benefit within three growing seasons due to increased

crop yields and reduced fertilizer costs. The scalability and affordability of graphene oxide and CNTs are still areas of active research. Recent advances in synthesis techniques, such as hydrothermal carbonization, are making these materials more economically viable for agricultural applications. Furthermore, the reduction in fertilizer and irrigation costs associated with CBNMs could offset their higher initial investment over time, particularly in water-scarce or nutrient-depleted regions.

9. POTENTIAL RISKS AND LIMITATIONS

While carbon-based nanomaterials (CBNMs) hold great promise for improving soil health and agricultural productivity, their widespread adoption faces a variety of risks and limitations (Biswas *et.al.*, 2024). These challenges include environmental concerns, technical and economic barriers, and regulatory and ethical issues. Addressing these challenges is crucial to ensuring the safe and effective use of CBNMs in agriculture.

9.1 Environmental Concerns

Despite their benefits, some CBNMs may pose toxicity risks to soil organisms, plants, and other components of the ecosystem. Studies have demonstrated that high concentrations of certain nanomaterials, such as carbon nanotubes (CNTs) and graphene oxide (GO), can negatively impact soil microbial communities and plant growth. For example, CNTs have been found to disrupt cellular functions in soil microbes by causing oxidative stress and damaging cell membranes, ultimately reducing microbial diversity and activity. Graphene oxide can interact with plant root systems in ways that impede growth. Studies have shown that excessive levels of graphene oxide in soil can lead to toxicity by altering water and nutrient uptake in plant roots or causing oxidative stress due to the generation of reactive oxygen species (ROS) (Zhang *et.al.*, 2020). These findings highlight the importance of determining safe concentration levels for the application of CBNMs to minimize unintended toxic effects. The bioaccumulation of CBNMs in plants and their potential transfer to higher trophic levels also raise concerns about food safety and ecosystem health. Although many studies report minimal risks at low concentrations, the long-term impacts of CBNM accumulation in soil ecosystems remain poorly understood and require further research. The potential leaching

of nanoparticles from soils into water bodies represents another significant environmental risk associated with CBNMs. Carbon-based nanomaterials, particularly those with high mobility in soil matrices, may migrate to surface water or groundwater systems, where they can pose risks to aquatic organisms and human health. CNTs and fullerenes have been shown to aggregate under certain environmental conditions, altering water chemistry and potentially causing toxic effects in aquatic ecosystems (Lawrence *et.al.*, 2016). The interaction of CBNMs with heavy metals or pesticides in soil can create complex chemical species, which may leach into water systems and exacerbate contamination risks. For example, biochar has been shown to adsorb contaminants such as arsenic and lead, but the stability of these adsorbed species over time remains uncertain. Leaching risks are particularly concerning in areas with high rainfall or irrigated agricultural systems, where water flow can transport nanoparticles over large distances. To mitigate these risks, it is essential to assess the mobility, persistence, and transformations of CBNMs in soils under various environmental conditions. Developing strategies to minimize leaching, such as functionalizing nanomaterials to enhance their stability in soils, could also reduce potential harm to water resources.

9.2 Technical and Economic Barriers

One of the most significant barriers to the widespread use of CBNMs in agriculture is their high cost of production. Advanced synthesis techniques for materials such as graphene, carbon nanotubes, and fullerenes, including chemical vapor deposition (CVD) and arc discharge methods, are expensive and require specialized equipment (Manawi *et.al.*, 2018). Although biochar is relatively cost-effective compared to other CBNMs, large-scale production of high-quality biochar with specific properties suitable for soil amendment can still be cost-prohibitive for many farmers, especially in low-income regions. Scalability remains another critical issue. Producing sufficient quantities of CBNMs to meet the demands of large-scale agricultural applications presents logistical challenges. Current production methods for high-purity nanomaterials are often energy-intensive and have low yields, limiting their feasibility for widespread adoption. Addressing these limitations will require the development of more cost-effective, energy-efficient, and scalable manufacturing processes, such as

hydrothermal or microwave-assisted methods for producing graphene oxide and biochar. The uniform application of CBNMs in agricultural soils poses another significant technical challenge. Due to their nanoscale properties, CBNMs tend to aggregate or disperse unevenly in soil matrices, which can lead to inconsistent effects on soil properties and plant growth. For example, uneven distribution of biochar in soil can create "hotspots" of nutrient availability, while other areas remain nutrient-deficient, reducing overall effectiveness. Monitoring the fate and behavior of CBNMs in soils is technically complex and requires advanced analytical tools, such as transmission electron microscopy (TEM) or spectroscopy techniques, to track their transformations and interactions (Roupcova *et.al.*, 2020). The lack of standardized methods for applying and monitoring CBNMs in agricultural settings further complicates their practical use. Developing user-friendly application protocols and reliable monitoring techniques is essential to address these challenges.

9.3 Regulation and Ethical Concerns

The lack of standardization in the production, characterization, and application of CBNMs presents a major regulatory challenge. Variations in the size, surface chemistry, and functionalization of nanomaterials can result in highly variable effects on soil health, making it difficult to establish universal guidelines for their safe and effective use. Regulatory agencies have yet to develop comprehensive frameworks for the assessment and approval of CBNMs in agriculture. The absence of clear regulations regarding the environmental and health risks of CBNMs limits their adoption and creates uncertainty for stakeholders (Virto *et.al.*, 2022). International collaboration between researchers, policymakers, and industry leaders is needed to establish standardized protocols for testing the safety and efficacy of CBNMs before their commercial deployment. Public perception of nanotechnology in agriculture remains a significant barrier to the widespread use of CBNMs. Concerns about the potential environmental and health risks associated with nanomaterials have led to skepticism and resistance among consumers and advocacy groups. Misinformation about nanotechnology and its applications has further contributed to public distrust, highlighting the need for effective science communication and education. Ethical concerns about the potential monopolization of

nanotechnology by large agribusinesses also raise questions about the accessibility of CBNMs for smallholder farmers. Ensuring that these technologies are affordable and accessible to all stakeholders, including low-income communities, will be crucial for fostering public acceptance and maximizing their benefits (Bhanye *et.al.*, 2024). To address these concerns, it is essential to engage the public in discussions about the benefits and risks of nanotechnology, promote transparency in research and development, and implement equitable policies that prioritize sustainability and social inclusion.

10. CURRENT STATE OF RESEARCH ON CBNMs IN AGRICULTURE

Carbon-based nanomaterials (CBNMs) have emerged as transformative tools in sustainable agriculture, offering the potential to improve soil health, enhance crop productivity, and mitigate environmental challenges. The research on CBNMs in agriculture is still evolving, with several recent advances and key findings highlighting their potential, as well as critical gaps that need to be addressed.

10.1 Recent Advances

Laboratory-scale experiments have provided foundational insights into the mechanisms by which CBNMs improve soil health and agricultural productivity. These controlled studies have demonstrated that materials such as biochar, graphene oxide (GO), and carbon nanotubes (CNTs) positively influence soil structure, nutrient availability, and microbial activity. For example, biochar has been shown to enhance soil cation exchange capacity (CEC), improve water retention, and reduce nutrient leaching in a variety of soils (Dey *et.al.*, 2023). Graphene oxide has been found to increase water-holding capacity and promote the slow release of nutrients, making it a promising tool for drought-prone regions. Laboratory studies have also highlighted the antimicrobial properties of certain CBNMs, such as graphene oxide and CNTs, which can suppress soil pathogens while supporting beneficial microbial communities. CNTs, in particular, have demonstrated the ability to improve plant nutrient uptake and seed germination in hydroponic systems and soil-based studies. A study showed that CNTs enhance the growth of tomato plants by facilitating water and nutrient transport to roots. These studies have laid the groundwork for understanding the effects of CBNMs on soil and

plants, setting the stage for field-scale applications. While lab-scale studies have highlighted the potential benefits of CBNMs, field trials and large-scale experiments are essential for evaluating their performance under real-world agricultural conditions. Recent field studies have begun to explore the use of biochar and graphene-based materials in crop production, demonstrating their potential to enhance soil fertility, water retention, and crop yields (Bhattacharya *et.al.*, 2023). A field study conducted in India demonstrated that biochar application increased soil organic carbon content and improved the water-holding capacity of degraded soils, resulting in a 20% increase in wheat yield. Another study in China evaluated the effects of graphene oxide on rice production, finding that GO improved water-use efficiency and nutrient uptake, leading to a 15% increase in grain yield. Field trials involving advanced CBNMs, such as CNTs and fullerenes, remain limited due to high costs and concerns about environmental safety. The scalability and economic feasibility of these materials are critical challenges that need to be addressed before their widespread adoption in agriculture.

10.2 Key Findings from Recent Studies

A growing body of research has demonstrated that CBNMs can significantly improve soil physical and chemical properties. For example, biochar's porous structure enhances soil aeration, water retention, and aggregation, making it particularly beneficial for sandy or degraded soils (Amoakwah *et.al.*, 2017). Studies have also shown that biochar application increases soil pH in acidic soils, improving nutrient availability for crops. Graphene oxide has been found to improve the soil's water-holding capacity and reduce the evaporation rate, which is crucial for water-scarce regions. Graphene oxide and biochar have demonstrated the ability to adsorb heavy metals and organic pollutants, reducing their bioavailability and toxicity in contaminated soils. These findings suggest that CBNMs can play a critical role in soil remediation and sustainable land management. CBNMs have shown significant positive effects on plant growth and crop yield in both lab and field studies. Biochar, has been consistently shown to enhance crop productivity by improving nutrient availability, water retention, and microbial activity in the rhizosphere. Graphene oxide and CNTs, due to their nanoscale properties, directly interact with plant roots, enhancing nutrient uptake and promoting

root elongation. A study found that CNTs enhanced the uptake of essential nutrients, such as nitrogen and potassium, resulting in increased biomass production in wheat plants. Graphene oxide has been shown to stimulate seed germination and improve the quality of rice grains (Li *et.al.*, 2018). These studies highlight the potential of CBNMs to improve crop productivity, particularly in regions facing soil degradation or water scarcity. The long-term impacts of these materials on plant health and food safety remain an important area of investigation.

10.3 Gaps in Current Research

While short-term studies have demonstrated the benefits of CBNMs, their long-term impacts on soil ecosystems and the environment remain poorly understood. Questions about the persistence of CBNMs in soils, their degradation pathways, and their interactions with soil organic matter and minerals require further investigation. For example, biochar is known for its long-term stability in soils, but its effects on soil microbial communities and nutrient cycling over decades are not fully understood. Graphene-based materials and CNTs may persist in soils for extended periods, potentially affecting microbial diversity and soil health in unforeseen ways (Das *et.al.*, 2023). The potential accumulation of CBNMs in soil ecosystems also raises concerns about their environmental fate and potential ecotoxicity. More research is needed to assess the impacts of these materials under varying environmental conditions and to establish safe application thresholds. Most studies on CBNMs have been conducted in controlled environments using specific soil types, such as sandy or loamy soils. The behavior and efficacy of CBNMs can vary significantly across different soil types, including clay, silt, and saline soils. The adsorption capacity of biochar and graphene oxide depends on soil texture, pH, and organic matter content. In clay-rich soils, CBNMs may exhibit reduced mobility and limited interaction with plant roots, whereas in sandy soils, they may enhance water retention and nutrient availability more effectively (Mukherjee *et.al.*, 2016). Understanding how CBNMs interact with diverse soil types is critical for optimizing their application in different agroecological zones. There is a lack of research on the effects of CBNMs in highly degraded or contaminated soils, where their potential to aid in soil restoration could be particularly valuable. Future studies should aim to explore the behavior and

impacts of CBNMs across a wider range of soil types and environmental conditions.

11. FUTURE

As the adoption of carbon-based nanomaterials (CBNMs) in agriculture expands, there is a growing need to address the technological, environmental, and policy challenges that accompany their use.

11.1 Research Priorities

One of the biggest barriers to the widespread use of CBNMs in agriculture is the high cost of production. Advanced nanomaterials such as carbon nanotubes (CNTs), graphene oxide (GO), and fullerenes require energy-intensive and complex synthesis techniques, such as chemical vapor deposition (CVD) or arc discharge, which limit scalability. Future research must prioritize the development of cost-effective and environmentally friendly methods for producing CBNMs at a large scale (Asghar *et.al.*, 2024). Emerging technologies, such as microwave-assisted synthesis and hydrothermal carbonization, have shown promise for reducing energy consumption and costs. For example, hydrothermal carbonization can convert agricultural waste into biochar or graphene-like materials under moderate temperature and pressure conditions, making it a sustainable option for nanomaterial production. Biological synthesis methods, such as using microorganisms or plant extracts to produce nanomaterials, represent a green and cost-effective alternative. Scaling up production will also require innovations in manufacturing processes to ensure consistent quality and functionalization of CBNMs. Research efforts should focus on optimizing material properties, such as porosity, surface area, and functional groups, to enhance their performance as soil amendments while minimizing production costs. The long-term impacts of CBNMs on soil health and ecosystem dynamics remain poorly understood. While short-term studies have demonstrated significant benefits, such as improved nutrient retention and plant growth, the persistence and transformation of CBNMs in soil environments require further investigation (Xiaobin *et.al.*, 2006). Future research should focus on understanding how CBNMs interact with soil components, such as organic matter, minerals, and microbial communities, over extended periods. Biochar is known to be stable in soils for decades to centuries, but its effects on nutrient

cycling and microbial activity may change over time. Graphene oxide and CNTs may undergo physical or chemical transformations that could affect their efficacy and potential toxicity. The environmental fate of CBNMs, including their potential to leach into water systems or accumulate in soils, needs to be studied under various environmental conditions. Long-term field experiments across diverse soil types and climates are essential to assess the sustainability and safety of CBNMs in agriculture (Biswas *et.al.*, 2024). These studies should also evaluate the impacts of CBNMs on soil biodiversity and ecosystem services, such as carbon sequestration and water filtration.

11.2 Integration with Other Technologies

The integration of CBNMs with precision agriculture and Internet of Things (IoT)-based systems represents a transformative approach to modern farming. Precision agriculture uses data-driven technologies, such as remote sensing, drones, and soil sensors, to optimize resource use and improve crop yields. By combining CBNMs with these technologies, farmers can achieve targeted soil amendments and enhanced monitoring capabilities. For example, graphene oxide-based sensors can be embedded in soils to monitor parameters such as nutrient levels, moisture content, and pH in real time. These sensors provide actionable insights, enabling precise application of fertilizers and irrigation, which minimizes waste and reduces environmental impacts. CNTs have shown potential in the development of advanced biosensors for detecting soil pathogens and contaminants, offering new tools for soil health monitoring (Kundu *et.al.*, 2019). Integrating CBNMs with IoT systems can also facilitate the remote management of agricultural fields, enabling farmers to make informed decisions based on real-time data. This synergy between nanotechnology and digital agriculture could revolutionize soil management practices, improving efficiency and sustainability at scale. CBNMs have the potential to play a pivotal role in regenerative agriculture, a holistic approach that focuses on restoring soil health, increasing biodiversity, and enhancing ecosystem resilience. Biochar, in particular, is well-suited for regenerative practices due to its ability to improve soil structure, sequester carbon, and promote microbial diversity. Future research should explore the use of CBNMs in conjunction with other regenerative practices, such as cover

cropping, crop rotation, and reduced tillage. Combining biochar with organic amendments, such as compost or green manure, could further enhance soil fertility and carbon sequestration (Kimetu *et.al.*, 2010). Graphene oxide and CNTs could be used to deliver biostimulants or beneficial microbes to the rhizosphere, supporting plant health and resilience under stress conditions such as drought or salinity. CBNMs could also aid in the reclamation of degraded lands by stabilizing soil structure and immobilizing pollutants. This aligns with the goals of regenerative agriculture to restore degraded ecosystems and create sustainable food systems.

11.3 Policy and Regulation

The lack of standardized guidelines for the production, application, and disposal of CBNMs in agriculture remains a critical challenge. Variability in the size, structure, and functionalization of nanomaterials can result in differing environmental and health impacts, making it essential to establish clear regulations for their use (Lead *et.al.*, 2018). Future policies should focus on developing standardized testing protocols to evaluate the safety and efficacy of CBNMs under diverse agricultural conditions. These guidelines should address issues such as application rates, potential toxicity thresholds, and environmental fate. Regulatory agencies, such as the FAO and national governments, should collaborate with researchers and industry stakeholders to create frameworks that balance innovation with safety. Policies should promote transparency in the production and labeling of CBNMs to ensure that farmers and consumers are informed about their benefits and risks. This will help build public trust and support for the adoption of nanotechnology in agriculture. Addressing the complex challenges associated with CBNMs in agriculture requires interdisciplinary collaboration among scientists, policymakers, and industry leaders. Chemists, soil scientists, ecologists, and agricultural engineers must work together to develop sustainable solutions that maximize the benefits of CBNMs while minimizing risks. Public-private partnerships could accelerate research and development by providing funding for large-scale experiments and pilot projects. For example, collaborations between nanotechnology companies and agricultural organizations could focus on creating affordable and scalable CBNM-based solutions for smallholder farmers (Ham *et.al.*, 1998).

12. CONCLUSION

Carbon-based nanomaterials (CBNMs) represent a transformative innovation in sustainable agriculture, offering solutions to critical challenges such as soil degradation, nutrient inefficiency, and climate change impacts. Their unique physical, chemical, and biological properties improve soil structure, water retention, nutrient availability, and microbial activity, leading to enhanced crop productivity and resilience. Their widespread adoption requires addressing key challenges, including cost-effective synthesis, environmental safety, and long-term impacts on soil ecosystems. Standardized regulations, interdisciplinary collaboration, and integration with technologies like precision agriculture are essential to ensure their safe and effective use. While research gaps persist, advances in scalable production and long-term studies highlight the immense potential of CBNMs to revolutionize soil management and support global food security. With responsible innovation, CBNMs could play a pivotal role in achieving sustainable agricultural practices and environmental resilience.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Asghar, N., Hussain, A., Nguyen, D. A., Ali, S., Hussain, I., Junejo, A., & Ali, A. (2024). Advancement in nanomaterials for environmental pollutants remediation: A systematic review on bibliometrics analysis, material types, synthesis pathways, and related mechanisms. *Journal of Nanobiotechnology*, 22(1), 26.
- Asghar, N., Hussain, A., Nguyen, D. A., Ali, S., Hussain, I., Junejo, A., & Ali, A. (2024). Advancement in nanomaterials for environmental pollutants remediation: A systematic review on bibliometrics analysis, material types, synthesis

- pathways, and related mechanisms. *Journal of Nanobiotechnology*, 22(1), 26.
- Chahar, M., Khaturia, S., Singh, H. L., Solanki, V. S., Agarwal, N., Sahoo, D. K., ... & Patel, A. (2023). Recent advances in the effective removal of hazardous pollutants from wastewater by using nanomaterials—A review. *Frontiers in Environmental Science*, 11, 1226101.
- Chahar, M., Khaturia, S., Singh, H. L., Solanki, V. S., Agarwal, N., Sahoo, D. K., ... & Patel, A. (2023). Recent advances in the effective removal of hazardous pollutants from wastewater by using nanomaterials: A review. *Frontiers in Environmental Science*, 11, 1226101.
- Cui, M., Zeng, L., Qin, W., & Feng, J. (2020). Measures for reducing nitrate leaching in orchards: A review. *Environmental Pollution*, 263, 114553.
- Cui, M., Zeng, L., Qin, W., & Feng, J. (2020). Measures for reducing nitrate leaching in orchards: A review. *Environmental Pollution*, 263, 114553.
- Davidson, D., & Gu, F. X. (2012). Materials for sustained and controlled release of nutrients and molecules to support plant growth. *Journal of Agricultural and Food Chemistry*, 60(4), 870-876.
- Davidson, D., & Gu, F. X. (2012). Materials for sustained and controlled release of nutrients and molecules to support plant growth. *Journal of Agricultural and Food Chemistry*, 60(4), 870-876.
- Farhat, S., & Scott, C. D. (2006). Review of the arc process modeling for fullerene and nanotube production. *Journal of Nanoscience and Nanotechnology*, 6(5), 1189-1210.
- Farhat, S., & Scott, C. D. (2006). Review of the arc process modeling for fullerene and nanotube production. *Journal of Nanoscience and Nanotechnology*, 6(5), 1189-1210.
- Fischer, B. M., Manzoni, S., Morillas, L., Garcia, M., Johnson, M. S., & Lyon, S. W. (2019). Improving agricultural water use efficiency with biochar—A synthesis of biochar effects on water storage and fluxes across scales. *Science of the Total Environment*, 657, 853-862.
- Fischer, B. M., Manzoni, S., Morillas, L., Garcia, M., Johnson, M. S., & Lyon, S. W. (2019). Improving agricultural water use efficiency with biochar: A synthesis of biochar effects on water storage and fluxes across scales. *Science of the Total Environment*, 657, 853-862.
- Hussain, A., Rehman, F., Rafeeq, H., Waqas, M., Asghar, A., Afsheen, N., ... & Iqbal, H. M. (2022). In-situ, ex-situ, and nano-remediation strategies to treat polluted soil, water, and air—A review. *Chemosphere*, 289, 133252.
- Hussain, A., Rehman, F., Rafeeq, H., Waqas, M., Asghar, A., Afsheen, N., ... & Iqbal, H. M. (2022). In-situ, ex-situ, and nano-remediation strategies to treat polluted soil, water, and air: A review. *Chemosphere*, 289, 133252.
- Kabiri, S., Degryse, F., Tran, D. N., da Silva, R. C., McLaughlin, M. J., & Losic, D. (2017). Graphene oxide: A new carrier for slow release of plant micronutrients. *ACS Applied Materials & Interfaces*, 9(49), 43325-43335.
- Kabiri, S., Degryse, F., Tran, D. N., da Silva, R. C., McLaughlin, M. J., & Losic, D. (2017). Graphene oxide: A new carrier for slow release of plant micronutrients. *ACS Applied Materials & Interfaces*, 9(49), 43325-43335.
- Lehman, R. M., Cambardella, C. A., Stott, D. E., Acosta-Martinez, V., Manter, D. K., Buyer, J. S., ... & Karlen, D. L. (2015). Understanding and enhancing soil biological health: The solution for reversing soil degradation. *Sustainability*, 7(1), 988-1027.
- Lehman, R. M., Cambardella, C. A., Stott, D. E., Acosta-Martinez, V., Manter, D. K., Buyer, J. S., ... & Karlen, D. L. (2015). Understanding and enhancing soil biological health: The solution for reversing soil degradation. *Sustainability*, 7(1), 988-1027.
- Liu, R., & Lal, R. (2012). Nanoenhanced materials for reclamation of mine lands and other degraded soils: A review. *Journal of Nanotechnology*, 2012(1), 461468.
- Liu, R., & Lal, R. (2012). Nanoenhanced materials for reclamation of mine lands and other degraded soils: A review. *Journal of Nanotechnology*, 2012(1), 461468.
- Nakhli, S. A. A., Delkash, M., Bakhshayesh, B. E., & Kazemian, H. (2017). Application of zeolites for sustainable agriculture: A review on water and nutrient retention. *Water, Air, & Soil Pollution*, 228, 1-34.
- Nakhli, S. A. A., Delkash, M., Bakhshayesh, B. E., & Kazemian, H. (2017). Application of zeolites for sustainable agriculture: A

- review on water and nutrient retention. *Water, Air, & Soil Pollution*, 228, 1-34.
- Pan, Y., Liu, X., Zhang, W., Liu, Z., Zeng, G., Shao, B., ... & Chen, M. (2020). Advances in photocatalysis based on fullerene C60 and its derivatives: Properties, mechanism, synthesis, and applications. *Applied Catalysis B: Environmental*, 265, 118579.
- Pan, Y., Liu, X., Zhang, W., Liu, Z., Zeng, G., Shao, B., ... & Chen, M. (2020). Advances in photocatalysis based on fullerene C60 and its derivatives: Properties, mechanism, synthesis, and applications. *Applied Catalysis B: Environmental*, 265, 118579.
- Patle, P. N., Kadu, P. R., Gabhane, A. R., Pharande, A. L., Bhagat, A. P., Bhojar, S. M., ... & Rahangdale, M. K. (2019). Consequences provoked due to excess application of agrochemical on soil health deterioration—A review for sustainable agriculture. *Journal of Pharmacognosy and Phytochemistry*, 8(2S), 63-66.
- Patle, P. N., Kadu, P. R., Gabhane, A. R., Pharande, A. L., Bhagat, A. P., Bhojar, S. M., ... & Rahangdale, M. K. (2019). Consequences provoked due to excess application of agrochemical on soil health deterioration: A review for sustainable agriculture. *Journal of Pharmacognosy and Phytochemistry*, 8(2S), 63-66.
- Reddy, D. H. K., Vijayaraghavan, K., Kim, J. A., & Yun, Y. S. (2017). Valorisation of post-sorption materials: Opportunities, strategies, and challenges. *Advances in Colloid and Interface Science*, 242, 35-58.
- Reddy, D. H. K., Vijayaraghavan, K., Kim, J. A., & Yun, Y. S. (2017). Valorisation of post-sorption materials: Opportunities, strategies, and challenges. *Advances in Colloid and Interface Science*, 242, 35-58.
- Reed, M. S., & Stringer, L. C. (2016). *Land degradation, desertification and climate change: Anticipating, assessing and adapting to future change*. Routledge.
- Reed, M. S., & Stringer, L. C. (2016). *Land degradation, desertification and climate change: Anticipating, assessing and adapting to future change*. Routledge.
- Sheoran, K., Kaur, H., Siwal, S. S., Saini, A. K., Vo, D. V. N., & Thakur, V. K. (2022). Recent advances of carbon-based nanomaterials (CBNMs) for wastewater treatment: Synthesis and application. *Chemosphere*, 299, 134364.
- Sheoran, K., Kaur, H., Siwal, S. S., Saini, A. K., Vo, D. V. N., & Thakur, V. K. (2022). Recent advances of carbon-based nanomaterials (CBNMs) for wastewater treatment: Synthesis and application. *Chemosphere*, 299, 134364.
- Thompson, A. W., Marzec, R., & Burniske, G. (2022). Climate BufferNet: A gaming simulation linking biodiversity conservation and climate change adaptation with agricultural landscape planning. *Landscape Journal*, 41(1), 45-60.
- Thompson, A. W., Marzec, R., & Burniske, G. (2022). Climate BufferNet: A gaming simulation linking biodiversity conservation and climate change adaptation with agricultural landscape planning. *Landscape Journal*, 41(1), 45-60.
- Vithanage, M., Seneviratne, M., Ahmad, M., Sarkar, B., & Ok, Y. S. (2017). Contrasting effects of engineered carbon nanotubes on plants: A review. *Environmental Geochemistry and Health*, 39, 1421-1439.
- Vithanage, M., Seneviratne, M., Ahmad, M., Sarkar, B., & Ok, Y. S. (2017). Contrasting effects of engineered carbon nanotubes on plants: A review. *Environmental Geochemistry and Health*, 39, 1421-1439.
- Wahab, A., Muhammad, M., Ullah, S., Abdi, G., Shah, G. M., Zaman, W., & Ayaz, A. (2024). Agriculture and environmental management through nanotechnology: Eco-friendly nanomaterial synthesis for soil-plant systems, food safety, and sustainability. *Science of the Total Environment*, 171862.
- Wahab, A., Muhammad, M., Ullah, S., Abdi, G., Shah, G. M., Zaman, W., & Ayaz, A. (2024). Agriculture and environmental management through nanotechnology: Eco-friendly nanomaterial synthesis for soil-plant systems, food safety, and sustainability. *Science of the Total Environment*, 171862.
- Wang, Z. H., & Li, S. X. (2019). Nitrate N loss by leaching and surface runoff in agricultural land: A global issue (a review). *Advances in Agronomy*, 156, 159-217.
- Wang, Z. H., & Li, S. X. (2019). Nitrate N loss by leaching and surface runoff in agricultural land: A global issue (a review). *Advances in Agronomy*, 156, 159-217.
- Wu, T., Liu, Y., Zheng, T., Dai, Y., Li, Z., & Lin, D. (2023). Fe-based nanomaterials and plant growth promoting rhizobacteria synergistically degrade polychlorinated biphenyls by producing extracellular reactive oxygen species. *Environmental Science & Technology*, 57(34), 12771-12781.

- Wu, T., Liu, Y., Zheng, T., Dai, Y., Li, Z., & Lin, D. (2023). Fe-based nanomaterials and plant growth-promoting rhizobacteria synergistically degrade polychlorinated biphenyls by producing extracellular reactive oxygen species. *Environmental Science & Technology*, 57(34), 12771-12781.
- Zhang, X., Cao, H., Wang, H., Zhao, J., Gao, K., Qiao, J., ... & Ge, S. (2022). The effects of graphene-family nanomaterials on plant growth: A review. *Nanomaterials*, 12(6), 936.
- Zhang, X., Cao, H., Wang, H., Zhao, J., Gao, K., Qiao, J., ... & Ge, S. (2022). The effects of graphene-family nanomaterials on plant growth: A review. *Nanomaterials*, 12(6), 936.
- Zhang, Y., Wang, J., & Feng, Y. (2021). The effects of biochar addition on soil physicochemical properties: A review. *Catena*, 202, 105284.
- Zhang, Y., Wang, J., & Feng, Y. (2021). The effects of biochar addition on soil physicochemical properties: A review. *Catena*, 202, 105284.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129020>